
Flowchart for Developmental-Behavioral Surveillance & Screening (0 to 6 years) in Medical Home Settings

From Marks, K.P., Glascoe, F.P., and Macias M.M. [2011] Enhancing the algorithm for developmental-behavioral surveillance and screening in children 0 to 5 years.



AAP Screening Recommendations

(a) General (broad-band) developmental screen routinely at

9, 18 & 24 or 30 mo. plus at 4 years to measure "kindergarten readiness", plus as needed if at risk for a developmental delay

(b) SE screen routinely at 5 years plus as needed if at risk for SE problem (c) Autism-specific

screen routinely at 18 & 24 mo, plus as needed if at risk for ASD

(d) Consider family psychosocial screen (e.g., 0 to 1 mo or at the initial visit), plus routinely employ a maternal mood disorder screen

in the first year

(eg, 2, 4 and/or 6 mo) Other Key Recommendations

(e) Evidence-based interventions are recommended for component 5. which is flexible & can occur before & after component 6

(f) A system-wide program with centralized care coordination or a practice-based integrative model of care (eg, colocation with developmental or mental health specialists) is recommended.

Legend Box = process

or action step Diamond = decision Y = yes N = no(+) = concerning (+/-) = borderline (-) = typical AAP = American Academy of Pediatrics

ASD = Austim Spectrum Disorder ECSE = early childhood special education EI = early intervention

IDEA = Individuals with Disabilities **Education Act** SE= social-emotional ROR = Reach Out & Read

WCV= well-child visit